December 2020

No.14 Total:14



Aerospace Information Research Institute, Chinese Academy of Sciences Key laboratory of Digital Earth Science, Chinese Academy of Sciences Big Earth Data Science Engineering Project (CASEarth) Sino-UK Crop Pest and Disease Forecasting & Management Joint Laboratory Key Lab of Aviation Plant Protection, Ministry of Agriculture and Rural Affairs, P.R. China National Engineering Research Center for Agro-Ecological Big Data Analysis & Application **Report of Monitoring and Assessment of Desert Locust in Africa and Asia** *Early December 2020*

Desert Locust monitoring and loss assessment in Somalia

Overview

Integrated with multi-source Farth Observation data, e.g. meteorological data, field data, and remote sensing data (such as GF series in China, MODIS and Landsat series in US, Sentinel series in EU), and selfdeveloped models and algorithms for Desert and Locust monitoring forecasting, the research team constructed the 'Vegetation pests and diseases monitoring and forecasting which could release system', regularly thematical maps and reports on Desert Locust.

This report focuses on the dynamic updates of desert locust monitoring and loss assessment in Somalia from October to November, 2020. The results showed that from October to November 2020, desert locusts in Somalia were mainly distributed in the northwest and the middle of the border with Ethiopia. Compared with September, the newly damaged vegetation area was 975.5 thousand hectares, including 0.6 thousand hectares of cropland, 129.5 thousand hectares of

Content	
Overview	1
Monitoring and assessment of Desert	
Locust in Somalia	2
Contact us	5

grassland, and 845.4 thousand hectares of the shrub. In December, the desert locust in Somalia will continue to reproduce and migrate southward. It is expected to reach northeastern Kenya in mid-December and spread westward and southward. The next three months coincides with the important crop growing season in Somalia. If not properly controlled, locusts will bring a major threat to agricultural and pasture production. It is necessary to continue the monitoring and early warning of the intercontinental desert locust plague, and organize joint prevention and control in multiple countries, ensuring the safety of agricultural and pasture production, as well as regional stability.

Monitoring and assessment of Desert Locust in Somalia

In early October. Yemeni locust swarms continued to move across the Gulf of Aden to northern Somalia. Somali locust swarms were mainly located in the Garowe region in the northwest and northeast. In mid-to-late October. the northeast locust swarms continued to multiply. Along with the north wind, the northern locust swarms gradually spread to the center and laid eggs, and the number of locust swarms in central Somalia continued to increase. The results showed that, by the end of October 2020, desert locust in Somalia harmed about 591.8 thousand hectares of vegetation area, an increase of 416.3 thousand hectares compared with September 2020 (including 0.1 thousand hectares of cropland, 47.7 thousand hectares of grassland, and 368.5 thousand hectares of shrub) (Figure 1). In early and mid-November, locusts in central Somalia continued to hatch, and the number of locusts continued to increase. Some locust swarms migrated south to northeastern and southern Kenya; in the middle and late November, with the heavy rainfall brought by tropical cyclone Gati, the locusts in the Garowe area continued to multiply and mature, leading to a further increase of the number of locusts. Coupled with the influence of the northerly winds, the locusts in central Somalia continued to spread to the south and eastern Kenya; during the same period, the locusts in eastern Ethiopia also continued to migrate eastward to northeast Somalia. The monitoring results showed that, by the end of November 2020, desert locust in Somalia harmed about 663.8 thousand hectares of vegetation area, an

increase of 559.2thousand hectares compared with September 2020 (including 0.5 thousand hectares of cropland, 81.8 thousand hectares of grassland, and 476.9 thousand hectares of shrub) (Figure 2).

The results showed that, from October to November 2020, desert locust in Somalia newly harmed about a total of 975.5 thousand hectares of vegetation area, including 0.6 thousand hectares of cropland, 129.5 thousand hectares of grassland, and 845.4 thousand hectares of shrub, accounting for 0.6%, 3.3% and 1.9% of the total cropland, grassland, and shrub in Somalia, respectively. The affected areas were mainly located in the northwest, central and southern parts of Somalia. Gedo in the south was the largest affected area (with affected area of 210.2 thousand hectares); followed by Hiiraan in the central (with affected area of 190.2 thousand hectares), and again Galguduud in the middle, Mudug in the middle, Bakool in the south, Wogooyi galbeed in the northwest, and Awdal in the northwest, and the affected areas were 141.4, 110.2, 69.8, 69.5, 59.4 thousand hectares respectively. Togdheer in the northwest had affected area of 38.8 thousand hectares, Sool in the north had affected area of 34 thousand hectares, Sanaag in the north had affected area of 22 thousand hectares, and Bay in the south had affected area of 15.5 thousand hectares; Bari in the north. Shabeellaha dhexe in the south, Nugaal in the middle and Jubbada dhexe in the south had less affected areas, with damaged area of 4.6, 1.8. 0.8 thousand 6.8, hectares. respectively.

This study used EU Sentinel-2 remote sensing images to monitor desert locust disasters in the more severely affected areas at the junction of Somalia and Ethiopia (Figure 3). The data acquisition time is October 2020, and the spatial resolution is 10m. The study area is located in the southern region of Odal, about 25 kilometers southeast of Boorama and 20 kilometers northeast of Weeraar. The vegetation types include grassland, shrubs and cropland, including 1.7 thousand hectares of cropland, 0.3 thousand hectares of grassland, and 180.3 thousand hectares of shrubland. The monitoring results showed that, in October 2020, the affected area of vegetation in the study area was 20.6 thousand hectares, accounting for 11.3% of the total area of the study area. Among them, the shrub was affected most severely with 20.1 thousand hectares, while the affected areas of cropland and grassland are 0.4 thousand hectares and

0.1 thousand hectares, accounting for 11.1%, 23.5%, and 33.3% of the total area of shrub, cropland and grassland, respectively. The results of the study show that desert locusts still threaten the vegetation in Somalia, and continuous monitoring of the locust situation is needed to ensure the agricultural production and food security in Somalia.

Comprehensive analysis shows that in December 2020, the locusts on the border between central Somalia and Ethiopia will continue to multiply and spread. It is expected that in mid-December, a large number of immature locusts will continue to migrate south, reach southern Somalia and invade northeastern Kenya, and spreading to the north and central counties.



Figure 1 Monitoring of Desert Locust damage in Somalia (October 2020)



Figure 2 Monitoring of Desert Locust damage in Somalia (November 2020)



Figure 3 Monitoring of Desert Locust damage in the key damage area of Somalia based on Sentinel-2 images (October 2020)

Contact us

Aerospace Information Research Institute Chinese Academy of Sciences No.9 Dengzhuang South Road, Haidian District, Beijing 100094, P.R.China. http://www.rscrop.com/ http://www.rscropmap.com





Chinese English The Vegetation Pests and Diseases Monitoring and Forecasting system are available under: http://www.rscropmap.com/

Legal Notice

Neither the Aerospace Information Research Institute nor any person action on behalf of the institute is responsible for the use which might be made of the publication.

Disclaimer

This report is a product of the Vegetation Remote Sensing & Pest and Disease Application Research Team of the Aerospace Information Research Institute, Chinese Academy of Sciences. The analyses and conclusions in the report do not represent the views of the Chinese Academy of Sciences or the Aerospace Information Research Institute. Users can legally quote the data in this report and indicate the source. However, any judgments, inferences or opinions made based on the report do not represent the views of the Team. The data published in this report are for reference only. The Team does not bear any legal responsibility arising from the use of the report. Official Chinese boundaries are used in the report.

Mission statements: As the science and knowledge service, the Sino-UK Crop Pest and Disease Forecasting & Management Joint Laboratory is to support independent evidence for crop monitoring.

Supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (XDA19080304), National Key R&D Program of China (2017YFE0122400, 2016YFB0501501), National Natural Science Foundation of China (61661136004, 41801338, 41801352, 41871339), Beijing Nova Program of Science and Technology (Z191100001119089), National special support program for high-level personnel recruitment (Wenjiang Huang), and Youth

Innovation Promotion Association CAS (2017085). Citation

Report of Monitoring and Assessment of Desert Locust in Africa and Asia, (2020). Desert Locust monitoring and loss assessment in Somalia. Beijing, China: RSCROP. DOI: 10.12237/casearth.5fd97934819aec0e3cddfd05.

Contact us Email: rscrop@aircas.ac.cn

Corresponding author

Professor Wenjiang Huang Aerospace Information Research Institute, Chinese Academy of Sciences Email: huanwj@aircas.ac.cn Tel: +86-10-82178178 FAX: +86-10-82178177

Main contributors

This report was released by Professor Wenjiang Huang's and Associate Professor Yingying Dong's research team in Aerospace Information Research Institute, Chinese Academy of Sciences.

Chinese contributors: Wenjiang Huang, Yingying Dong, Longlong Zhao, Huichun Ye, Mingquan Wu, Kun Wang, Xiaoping Du, Changyong Dou, Jun Yan, Jingcheng Zhang, Bei Cui, Linsheng Huang, Dailiang Peng, Hong Chang, Yun Geng, Chao Ruan, Huiqin Ma, Anting Guo, Linyi Liu, Naichen Xing, Yue Shi, Qiong Zheng, Yu Ren, Hansu Zhang, Tingguang Hu, Yanru Huang, Yu Jin, Chao Ding, Biyao Zhang, Zhongxiang Sun, Xiangmei Qin, Xueling Li, , Yingxin Xiao, Zhuoqing Hao, Kang Wu, Yong Liu, Bo Wu, Weiping Kong, Juhua Luo, Jinling Zhao, Dongyan Zhang, Xiaodong Yang, Yanhua Meng, Wenjie Fan, Yue Liu, Gang Sun, Bin Wu, Qing Zhang, Dacheng Wang, Wei Feng, Xianfeng Zhou, Qiaoyun Xie, Muyi Huang, Jing Jiang, Zhaochuan Wu, Cuicui Tang, Fang Xu, Jianli Li, Wenjing Liu, Junjing Lu, Furan Song, Qingsong Guan, Qinying Yang, Chuang Liu.

Foreign contributors: Belinda Luke, Bethan Perkins, Bryony Taylor, Hongmei Li, Wenhua Chen, Pablo Gonzalez-Moreno, Sarah Thomas, Timothy Holmes, Stefano Pignatti, Giovanni Laneve, Raffaele Casa, Simone Pascucci, Martin Wooster, Jason Chapman.

Advisory Experts: Bing Zhang, Gensuo Jia, Jihua Wang, Qiming Qin, Puyun Yang, Guofei Fang, Shouquan Chai, Jingquan Zhu, Yuying Jiang, Zhonghua Zhao, Binyuan Ren, Dongmei Yan, Xiangtao Fan, Jianhui Li, Jie Liu, Yubin Lan, Jingfeng Huang, Anhong Guo, Zhanhong Ma, Yilin Zhou, Xiongbing Tu, Wenbing Wu, Feng Zhang, Zhiguo Wang, Lifang Wu, Dong Liang, Yanbo Huang, Chenghai Yang, Liangxiu Han, Ruiliang Pu, Hugh Mortimer, Jon Styles, Andy Shaw, Jadu Dash.